

**Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application.

1-123. (canceled).

124. (previously presented) An isolated nucleic acid comprising:

- (a) the nucleic acid sequence of SEQ ID NO: 348;
- (b) the full-length coding sequence of the nucleic acid sequence of SEQ ID NO: 348;

or

(c) the full-length coding sequence of the cDNA deposited under ATCC accession number 203044.

125-128. (canceled)

129. (previously presented) The isolated nucleic acid of Claim 124 comprising the nucleic acid sequence of SEQ ID NO: 348.

130. (previously presented) The isolated nucleic acid of Claim 124 comprising the full-length coding sequence of the nucleic acid sequence of SEQ ID NO: 348.

131. (previously presented) The isolated nucleic acid of Claim 124 comprising the full-length coding sequence of the cDNA deposited under ATCC accession number 203044.

132-134. (canceled)

135. (previously presented) A vector comprising the nucleic acid of Claim 124.

136. (previously presented) The vector of Claim 135, wherein said nucleic acid is operably linked to control sequences recognized by a host cell transformed with the vector.

137. (previously presented) A host cell comprising the vector of Claim 135.

138. (previously presented) The host cell of Claim 137, wherein said cell is a CHO cell, an *E. coli* or a yeast cell.

139. (currently amended) An isolated nucleic acid molecule consisting of a fragment of the nucleic acid sequence of SEQ ID NO: 348, or a complement thereof, of at least 20 nucleotides in length that hybridizes under stringent conditions to:

- (a) the nucleic acid sequence of SEQ ID NO: 348 or a complement thereof;
- (b) the full-length coding sequence of the cDNA deposited under ATCC accession number 203044 or a complement thereof;

wherein, said stringent conditions use 50% formamide, 5X SSC, 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5X Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, and washes at 42°C in 0.2X SSC, at 55°C in 50% formamide followed by a high-stringency wash at 55°C in 0.1X SSC, EDTA; wherein said isolated nucleic acid molecule is suitable for PCR use as a PCR primer or probe.

140. (previously presented) The isolated nucleic acid molecule of Claim 139 that is at least 50 nucleotides or above in length.

141. (previously presented) The isolated nucleic acid molecule of Claim 139 that is at least 60 nucleotides or above in length.

142. (previously presented) The isolated nucleic acid molecule of Claim 139 that is at least 70 nucleotides or above in length.

143. (previously presented) The isolated nucleic acid molecule of Claim 139 that is at least 80 nucleotides or above in length.

144. (previously presented) The isolated nucleic acid molecule of Claim 139 that is at least 90 nucleotides or above in length.

145. (previously presented) The isolated nucleic acid molecule of Claim 139 that is at least 100 nucleotides or above in length.